

Variational Inference with Implicit and Semi-Implicit Distributions

Francisco J. R. Ruiz

Marie Skłodowska-Curie Fellow


Deep|Bayes Summer School
August 24, 2019



This research is supported by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 706760

Notation

- ▶ Model: Joint distribution $p(x, z)$
- ▶ Latent variables z
- ▶ Observations x



The Posterior Distribution

$$p(z | x) = \frac{p(x, z)}{\int p(x, z) dz}$$

- ▶ The posterior allows us to explore the data and make predictions
- ▶ Intractable in general
- ▶ Approximate the posterior: Bayesian inference

Variational Inference (Quick Review)


$$p(z | x) = \frac{p(x, z)}{\int p(x, z) dz}$$

- ▶ Define a simple family of distributions $q_\theta(z)$ with parameters θ
- ▶ Fit θ by minimizing the KL divergence to the posterior,


$$\theta^* = \arg \min_{\theta} \text{KL}(q_\theta(z) || p(z | x))$$

- ▶ Variational inference solves an optimization problem

Variational Inference (Quick Review)



Variational Inference (Quick Review)



Variational Inference (Quick Review)

- ▶ Minimizing the KL \equiv Maximizing the ELBO

$$\mathcal{L}(\theta) = \mathbb{E}_{q_\theta(z)} [\log p(x, z) - \log q_\theta(z)]$$


- ▶ Variational inference finds θ to maximize $\mathcal{L}(\theta)$

Mean-Field Variational Inference


- ▶ Classical VI: Mean-field variational distribution:

$$q_{\theta}(z) = \prod_n q_{\theta_n}(z_n)$$

- ▶ Useful, simple, and fast, but might not be accurate



This Lecture: Expand the Variational Family



Beyond the Mean-Field Family

- ▶ **Structured VI** [Saul+, 1996; Ghahramani+, 1997; Titsias+, 2011]
- ▶ **Mixtures** [Bishop+, 1998; Gershman+, 2012; Salimans+, 2013; Guo+, 2016; Miller+, 2017]
- ▶ **Sampling mechanisms** [Salimans+, 2015; Hoffman, 2017; Maddison+, 2017; Naesseth+, 2017; Li+, 2017; Titsias, 2017; Naesseth+, 2018; Le+, 2018; Grover+, 2018; Zhang+, 2018; Habib+, 2019; Neklyudov+, 2019; Ruiz+, 2019]
- ▶ **Spectral methods** [Shi+, 2018]
- ▶ **Linear response estimates** [Giordano+, 2015; Giordano+, 2017]
- ▶ **Copulas** [Tran+, 2015; Han+, 2016]
- ▶ **Invertible transformations** [Rezende+, 2014; Kingma+, 2014; Titsias+, 2014; Kucukelbir+, 2015] & **Normalizing flows** [Rezende+, 2015; Kingma+, 2016; Papamakarios+, 2017; Tomczak+, 2016; Tomczak+, 2017; Dinh+, 2017]
- ▶ **Hierarchical models** [Ranganath+, 2016; Tran+, 2016; Maaløe+, 2016; Sobolev+, 2019]
- ▶ **Implicit distributions** [Mescheder+, 2017; Huszár, 2017; Tran+, 2017; Shi+, 2018] & **Semi-implicit distributions** [Yin+, 2018; Titsias+, 2019; Molchanov+, 2019]

Beyond the Mean-Field Family


- ▶ **Structured VI** [Saul+, 1996; Ghahramani+, 1997; Titsias+, 2011]
- ▶ **Mixtures** [Bishop+, 1998; Gershman+, 2012; Salimans+, 2013; Guo+, 2016; Miller+, 2017]
- ▶ **Sampling mechanisms** [Salimans+, 2015; Hoffman, 2017; Maddison+, 2017; Naesseth+, 2017; Li+, 2017; Titsias, 2017; Naesseth+, 2018; Le+, 2018; Grover+, 2018; Zhang+, 2018; Habib+, 2019; Neklyudov+, 2019; Ruiz+, 2019]
- ▶ **Spectral methods** [Shi+, 2018]
- ▶ **Linear response estimates** [Giordano+, 2015; Giordano+, 2017]
- ▶ **Copulas** [Tran+, 2015; Han+, 2016]
- ▶ **Invertible transformations** [Rezende+, 2014; Kingma+, 2014; Titsias+, 2014; Kucukelbir+, 2015] & **Normalizing flows** [Rezende+, 2015; Kingma+, 2016; Papamakarios+, 2017; Tomczak+, 2016; Tomczak+, 2017; Dinh+, 2017]
- ▶ **Hierarchical models** [Ranganath+, 2016; Tran+, 2016; Maaløe+, 2016; Sobolev+, 2019]
- ▶ **Implicit distributions** [Mescheder+, 2017; Huszár, 2017; Tran+, 2017; Shi+, 2018] & **Semi-implicit distributions** [Yin+, 2018; Titsias+, 2019; Molchanov+, 2019]

This Lecture

- ▶ Expand the variational family $q_\theta(z)$
- ▶ Use *implicit distributions*
 - ▶ Easy to sample from, $z \sim q_\theta(z)$
 - ▶ Intractable density, $q_\theta(z)$
- ▶ Challenge: Solve the optimization problem with intractable $q_\theta(z)$

objective: $\mathcal{L}(\theta) = \mathbb{E}_{q_\theta(z)} [\log p(x, z) - \log q_\theta(z)]$


Goal: More Expressive Variational Distributions



Part I:


Implicit Distributions and Adversarial Training

How to Form an Expressive Implicit Distribution




- ▶ Generate random noise $\varepsilon \sim q(\varepsilon)$
- ▶ Pass the noise through a NN with parameters θ
- ▶ Let z be the output of the NN

How to Form an Expressive Implicit Distribution



- ▶ Implicit distribution $q_\theta(z)$:
 - ▶ Easy to draw samples:
$$\text{sample } \varepsilon \sim q(\varepsilon); \quad \text{set } z = f_\theta(\varepsilon)$$
 - ▶ Cannot evaluate the density $q_\theta(z)$
- ▶ Flexible distribution $q_\theta(z)$ due to the NN
- ▶ Goal: Tune θ so that $q_\theta(z)$ approximates the posterior $p(z | x)$

Why VI with Implicit Distributions is Hard




- ▶ The VI objective is the ELBO (equivalent to minimizing KL),

$$\mathcal{L}(\theta) = \mathbb{E}_{q_\theta(z)} \left[\underbrace{\log p(x, z)}_{\text{model}} - \underbrace{\log q_\theta(z)}_{\text{entropy}} \right]$$

- ▶ Gradient of the objective $\nabla_\theta \mathcal{L}(\theta)$ (reparameterization)

$$\nabla_\theta \mathcal{L}(\theta) = \mathbb{E}_{q(\varepsilon)} \left[\nabla_\theta \left(\log p(x, f_\theta(\varepsilon)) - \log q_\theta(f_\theta(\varepsilon)) \right) \right]$$

Why VI with Implicit Distributions is Hard




- ▶ Gradient of the objective $\nabla_{\theta} \mathcal{L}(\theta)$ (reparameterization)

$$\nabla_{\theta} \mathcal{L}(\theta) = \mathbb{E}_{q(\varepsilon)} \left[\nabla_{\theta} \left(\log p(x, f_{\theta}(\varepsilon)) - \log q_{\theta}(f_{\theta}(\varepsilon)) \right) \right]$$

- ▶ For the model term:

$$\mathbb{E}_{q(\varepsilon)} [\nabla_{\theta} \log p(x, f_{\theta}(\varepsilon))] \approx \frac{1}{S} \sum_{s=1}^S \nabla_{\theta} \log p(x, f_{\theta}(\varepsilon^{(s)})), \quad \varepsilon^{(s)} \sim q(\varepsilon)$$

Why VI with Implicit Distributions is Hard



- ▶ Gradient of the objective $\nabla_{\theta} \mathcal{L}(\theta)$ (reparameterization)


$$\nabla_{\theta} \mathcal{L}(\theta) = \mathbb{E}_{q(\varepsilon)} \left[\nabla_{\theta} \left(\log p(x, f_{\theta}(\varepsilon)) - \log q_{\theta}(f_{\theta}(\varepsilon)) \right) \right]$$

- ▶ For the entropy term:

$$\nabla_{\theta} \log q_{\theta}(f_{\theta}(\varepsilon)) = \nabla_z \log q_{\theta}(z) \times \nabla_{\theta} f_{\theta}(\varepsilon) + \underbrace{\nabla_{\theta} \log q_{\theta}(z) \Big|_{z=f_{\theta}(\varepsilon)}}_{=0 \text{ (in expectation)}}$$

- ▶ Monte Carlo estimates require $\nabla_z \log q_{\theta}(z)$ (not available)

How Density Ratio Estimation Can Help




- ▶ The VI objective is the ELBO (equivalent to minimizing KL),

$$\mathcal{L}(\theta) = \mathbb{E}_{q_\theta(z)} \left[\underbrace{\log p(x, z)}_{\text{model}} - \underbrace{\log q_\theta(z)}_{\text{entropy}} \right]$$

- ▶ Rewrite the ELBO as “log-likelihood minus KL to prior,”

$$\begin{aligned}\mathcal{L}(\theta) &= \mathbb{E}_{q_\theta(z)} [\log p(x | z)] - \text{KL}(q_\theta(z) || p(z)) \\ &= \mathbb{E}_{q_\theta(z)} [\log p(x | z)] - \mathbb{E}_{q_\theta(z)} \left[\log \frac{q_\theta(z)}{p(z)} \right]\end{aligned}$$

How Density Ratio Estimation Can Help



- ▶ ELBO objective:

$$\mathcal{L}(\theta) = \mathbb{E}_{q_\theta(z)} [\log p(x | z)] - \mathbb{E}_{q_\theta(z)} \left[\log \frac{q_\theta(z)}{p(z)} \right]$$

- ▶ Key idea: Approximate the density ratio $\log \frac{q_\theta(z)}{p(z)}$

Density Ratio Estimation

- ▶ ELBO objective:

$$\mathcal{L}(\theta) = \mathbb{E}_{q_\theta(z)} [\log p(x | z)] - \mathbb{E}_{q_\theta(z)} \left[\log \frac{q_\theta(z)}{p(z)} \right]$$

- ▶ Imagine that we had labelled samples from $q_\theta(z)$ and $p(z)$:
 - Class $y = 1$: The sample z comes from $q_\theta(z)$
 - Class $y = 0$: The sample z comes from $p(z)$
- ▶ If you observe z , what is the class? (under equal class prior)
 - ▶ Optimal classifier is $D^*(z) = \frac{q_\theta(z)}{q_\theta(z) + p(z)}$
- ▶ The density ratio can be expressed as a function of the classifier:

$$\log \frac{q_\theta(z)}{p(z)} = \log D^*(z) - \log(1 - D^*(z))$$

Density Ratio Estimation

- ▶ The density ratio can be expressed as a function of the classifier:

$$\log \frac{q_\theta(z)}{p(z)} = \log D^*(z) - \log(1 - D^*(z))$$

- ▶ Train a (flexible) classifier $D(z)$ that distinguishes samples:

$$D^*(z) = \max_D \mathbb{E}_{q_\theta(z)} [D(z)] + \mathbb{E}_{p(z)} [\log(1 - D(z))]$$

- ▶ Rewrite the ELBO using $D(z)$,

$$\mathcal{L}(\theta) = \mathbb{E}_{q_\theta(z)} [\log p(x | z)] - \mathbb{E}_{q_\theta(z)} [\log D(z) - \log(1 - D(z))]$$

Density Ratio Estimation: Optimization

- ▶ ELBO objective:

$$\mathcal{L}(\theta) = \mathbb{E}_{q_\theta(z)} [\log p(x | z)] - \mathbb{E}_{q_\theta(z)} [\log D(z) - \log(1 - D(z))]$$

- ▶ Algorithm:

1. Follow gradient estimates of the ELBO w.r.t. θ (reparameterization)
2. For each θ , fit a flexible classifier $D(z)$ so that $D(z) \approx D^*(z)$

Limitations of Density Ratio Estimation

- ▶ ELBO objective:

$$\mathcal{L}(\theta) = \mathbb{E}_{q_\theta(z)} [\log p(x | z)] - \mathbb{E}_{q_\theta(z)} [\log D(z) - \log(1 - D(z))]$$

- ▶ Limitations:

- The discriminator $D(z)$ needs to be trained to optimum after each update of θ (in practice, optimization is truncated to a few iterations)
- Unstable training when discriminator does not catch up quickly
- In **high dimensions**, the discriminator overfits easily, giving values close to 0 or 1

Alternatives

- ▶ Kernel-based density ratio estimation (KIVI) [Shi+, 2018]
- ▶ Semi-implicit distributions [Yin+, 2018; Titsias+, 2019; Molchanov+, 2019]


Alternatives

- ▶ Kernel-based density ratio estimation (KIVI) [Shi+, 2018]
- ▶ Semi-implicit distributions [Yin+, 2018; Titsias+, 2019; Molchanov+, 2019]

Part II:

Semi-Implicit Distributions

Recap: VI with Implicit Distributions



- ▶ ELBO objective:

$$\mathcal{L}(\theta) = \mathbb{E}_{q_\theta(z)} \left[\underbrace{\log p(x, z)}_{\text{model}} - \underbrace{\log q_\theta(z)}_{\text{entropy}} \right]$$

- ▶ Gradient of the objective $\nabla_\theta \mathcal{L}(\theta)$ (reparameterization)

$$\nabla_\theta \mathcal{L}(\theta) = \mathbb{E}_{q(\varepsilon)} \left[\nabla_\theta \left(\log p(x, f_\theta(\varepsilon)) - \log q_\theta(f_\theta(\varepsilon)) \right) \right]$$

- ▶ Monte Carlo estimates require $\nabla_z \log q_\theta(z)$ (not available)

Semi-Implicit Distributions


- ▶ ELBO objective:

$$\mathcal{L}(\theta) = \mathbb{E}_{q_\theta(z)} \left[\underbrace{\log p(x, z)}_{\text{model}} - \underbrace{\log q_\theta(z)}_{\text{entropy}} \right]$$


- ▶ Goal: Tractable inference avoiding density ratio estimation
- ▶ Two methods:
 - Lower-bound the ELBO (SIVI) [Yin+, 2018; Molchanov+, 2019]
 - Estimate gradients with sampling (UIVI) [Titsias+, 2019]
- ▶ First step: use a semi-implicit construction of $q_\theta(z)$

Semi-Implicit Distributions

- ▶ Implicit distribution:




- ▶ (Semi-)implicit distribution:



Semi-Implicit Distributions

- ▶ (Semi-)implicit distribution




- ▶ **Example:** The conditional $q_\theta(z | \varepsilon)$ is a Gaussian,

$$q_\theta(z | \varepsilon) = \mathcal{N}(z | \mu_\theta(\varepsilon), \Sigma_\theta(\varepsilon))$$

Semi-Implicit Distributions

- ▶ (Semi-)implicit distribution



- ▶ The distribution $q_\theta(z)$ is still **implicit**,
 - ▶ Easy to sample,

sample $\varepsilon \sim q(\varepsilon)$,

obtain $\mu_\theta(\varepsilon)$ and $\Sigma_\theta(\varepsilon)$


sample $z \sim \mathcal{N}(z | \mu_\theta(\varepsilon), \Sigma_\theta(\varepsilon))$

- ▶ The variational distribution $q_\theta(z)$ is not tractable,

$$q_\theta(z) = \int q(\varepsilon)q_\theta(z | \varepsilon)d\varepsilon$$

Semi-Implicit Distributions


- ▶ (Semi-)implicit distribution



- ▶ **Assumptions** on the conditional $q_\theta(z | \varepsilon)$:
 - ▶ Reparameterizable
 - ▶ Tractable gradient $\nabla_z \log q_\theta(z | \varepsilon)$
Note: this is different from $\nabla_z \log q_\theta(z)$ (still intractable)

Semi-Implicit Distributions

- ▶ (Semi-)implicit distribution



- ▶ The Gaussian meets both assumptions:

- ▶ Reparameterizable,

$$u \sim \mathcal{N}(u | 0, I), \quad z = h_\theta(u ; \varepsilon) = \mu_\theta(\varepsilon) + \Sigma_\theta(\varepsilon)^{1/2} u$$

- ▶ Tractable gradient,

$$\nabla_z \log q_\theta(z | \varepsilon) = -\Sigma_\theta(\varepsilon)^{-1}(z - \mu_\theta(\varepsilon))$$

Method 1: SIVI

- ▶ Define a lower bound of the ELBO,

$$\mathcal{L}(\theta) \geq \bar{\mathcal{L}}(\theta), \quad \text{where}$$

$$\begin{aligned}\bar{\mathcal{L}}(\theta) = & \mathbb{E}_{\varepsilon \sim q(\varepsilon)} \left[\mathbb{E}_{z \sim q_\theta(z | \varepsilon)} \left[\mathbb{E}_{\varepsilon^{(1)}, \dots, \varepsilon^{(L)} \sim q(\varepsilon)} \left[\log p(x, z) \right. \right. \right. \\ & \left. \left. \left. - \log \left(\frac{1}{L+1} \left(q_\theta(z | \varepsilon) + \sum_{\ell=1}^L q_\theta(z | \varepsilon^{(\ell)}) \right) \right) \right] \right]\end{aligned}$$

- ▶ Optimize the lower bound instead of the ELBO
- ▶ The lower bound does not depend on the intractable $q_\theta(z)$

Method 1: SIVI

- ▶ SIVI bound:

$$\begin{aligned}\bar{\mathcal{L}}(\theta) = & \mathbb{E}_{\varepsilon \sim q(\varepsilon)} \left[\mathbb{E}_{z \sim q_\theta(z | \varepsilon)} \left[\mathbb{E}_{\varepsilon^{(1)}, \dots, \varepsilon^{(L)} \sim q(\varepsilon)} \left[\log p(x, z) \right. \right. \right. \\ & \left. \left. \left. - \log \left(\frac{1}{L+1} \left(q_\theta(z | \varepsilon) + \sum_{\ell=1}^L q_\theta(z | \varepsilon^{(\ell)}) \right) \right) \right] \right]\end{aligned}$$

- ▶ Free parameter L controls the tightness of the bound
 - As $L \rightarrow \infty$, $\bar{\mathcal{L}}(\theta) \rightarrow \mathcal{L}(\theta)$
 - Computational complexity increases with L
- ▶ SIVI allows for semi-implicit construction of prior in VAEs
[Molchanov+, 2019]

Method 2: UIVI


- ▶ Recall the reparameterization gradient of the ELBO,

$$\nabla_{\theta} \mathcal{L}(\theta) = \mathbb{E}_{q(\varepsilon)} \left[\nabla_{\theta} \left(\log p(x, f_{\theta}(\varepsilon)) - \log q_{\theta}(f_{\theta}(\varepsilon)) \right) \right]$$

- ▶ UIVI obtains an unbiased Monte Carlo estimator of $\nabla_z \log q_{\theta}(z)$
 - Avoid density ratio estimation
 - Directly optimize the ELBO (instead of a bound)
- ▶ Key idea: Gradient of the entropy component as an expectation,

$$\nabla_z \log q_{\theta}(z) = \mathbb{E}_{\text{distrib}(\cdot)} [\text{function}(z, \cdot)]$$

Method 2: UIVI




- ▶ Rewrite as an expectation,

$$\nabla_z \log q_\theta(z) = \mathbb{E}_{q_\theta(\varepsilon' | z)} [\nabla_z \log q_\theta(z | \varepsilon')]$$

- ▶ Form Monte Carlo estimate,

$$\nabla_z \log q_\theta(z) \approx \nabla_z \log q_\theta(z | \varepsilon'), \quad \varepsilon' \sim q_\theta(\varepsilon' | z)$$

Method 2: UIVI (Full Algorithm)




- ▶ The gradient of the ELBO is

$$\nabla_\theta \mathcal{L}(\theta) = \mathbb{E}_{q(\varepsilon)q(u)} \left[\nabla_z (\log p(x, z) - \log q_\theta(z)) \Big|_{z=h_\theta(u; \varepsilon)} \times \nabla_\theta h_\theta(u; \varepsilon) \right]$$

- ▶ Estimate the gradient based on samples:

1. Sample $\varepsilon \sim q(\varepsilon)$, $u \sim q(u)$ (standard Gaussians)
2. Set $z = h_\theta(\varepsilon; u) = \mu_\theta(\varepsilon) + \Sigma_\theta(\varepsilon)^{1/2} u$
3. Evaluate $\nabla_z \log p(x, z)$ and $\nabla_\theta h_\theta(u; \varepsilon)$
4. Sample $\varepsilon' \sim q_\theta(\varepsilon' | z)$
5. Approximate $\nabla_z \log q_\theta(z) \approx \nabla_z \log q_\theta(z | \varepsilon')$

Method 2: UIVI (The Reverse Conditional)




- ▶ The distribution $q_\theta(\varepsilon' | z)$ is the **reverse conditional**
The conditional is $q_\theta(z | \varepsilon)$
- ▶ Sample from $q_\theta(\varepsilon' | z)$ using HMC, targeting

$$q(\varepsilon' | z) \propto q(\varepsilon') q_\theta(z | \varepsilon')$$


- ▶ Problem: HMC is slow... How to accelerate this?

Method 2: UIVI (The Reverse Conditional)



- ▶ Recall the UIVI algorithm,
 1. Sample $\varepsilon \sim q(\varepsilon)$, $u \sim q(u)$ (standard Gaussians)
 2. Set $z = h_\theta(\varepsilon; u) = \mu_\theta(\varepsilon) + \Sigma_\theta(\varepsilon)^{1/2} u$
 3. Evaluate $\nabla_z \log p(x, z)$ and $\nabla_\theta h_\theta(u; \varepsilon)$
 4. Sample $\varepsilon' \sim q_\theta(\varepsilon' | z)$
 5. Approximate $\nabla_z \log q_\theta(z) \approx \nabla_z \log q_\theta(z | \varepsilon')$
- ▶ We have that $(\varepsilon, z) \sim q_\theta(\varepsilon, z) = q(\varepsilon)q_\theta(z | \varepsilon) = q_\theta(z)q_\theta(\varepsilon | z)$
- ▶ Thus, ε is a sample from $q_\theta(\varepsilon | z)$
- ▶ To accelerate sampling $\varepsilon' \sim q(\varepsilon' | z)$, initialize HMC at ε

Method 2: UIVI (The Reverse Conditional)




- ▶ Sample from $q_\theta(\varepsilon' | z)$ using HMC targeting

$$q(\varepsilon' | z) \propto q(\varepsilon') q_\theta(z | \varepsilon')$$



- ▶ Initialize HMC at stationarity (using ε)
- ▶ A few HMC iterations to reduce correlation between ε and ε'

UIVI: Toy Experiments



UIVI: Multinomial Logistic Regression Experiments

$$p(x, z) = p(z) \prod_{n=1}^N \frac{\exp\{x_n^\top z_{y_n} + z_0 y_n\}}{\sum_k \exp\{x_n^\top z_k + z_0 k\}}$$



UIVI provides better ELBO and predictive performance than SIVI

UIVI: VAE Experiments

- ▶ Model is $p_\phi(x, z) = \prod_n p(z_n)p_\phi(x_n | z_n)$
- ▶ Amortized variational distrib. $q_\theta(z_n | x_n) = \int q(\varepsilon_n) q_\theta(z_n | \varepsilon_n, x_n) d\varepsilon_n$
- ▶ Goal: Find model parameters ϕ and variational parameters θ


method	average test log-likelihood	
	MNIST	Fashion-MNIST
Explicit (standard VAE)	-98.29	-126.73
SIVI	-97.77	-121.53
UIVI	-94.09	-110.72

UIVI provides better predictive performance

Part III:

MCMC-Improved Approximation

Our Goal: More Expressive Variational Distributions



Main Idea: Use MCMC

- ▶ Start from an *explicit* variational distribution, $q_\theta^{(0)}(z)$
- ▶ Improve the distribution with t MCMC steps,

$$z_0 \sim q_\theta^{(0)}(z), \quad z \sim Q^{(t)}(z | z_0)$$

(the MCMC sampler targets the posterior, $p(z | x)$)

- ▶ Implicit variational distribution,

$$q_\theta(z) = \int q_\theta^{(0)}(z_0) Q^{(t)}(z | z_0) dz_0$$

Challenges of Using MCMC in VI

$$\mathcal{L}_{\text{improved}}(\theta) = \mathbb{E}_{q_\theta(z)} [\log p(x, z) - \log q_\theta(z)]$$

- ▶ Challenge #1: The variational objective becomes intractable
- ▶ Challenge #2: The variational objective may depend *weakly* on θ

$$q_\theta(z) \xrightarrow{t \rightarrow \infty} p(z | x)$$

Alternative Divergence: VCD

- ▶ We would like an objective that avoids these challenges
- ▶ We call the objective *Variational Contrastive Divergence*, $\mathcal{L}_{\text{VCD}}(\theta)$
- ▶ Desired properties:
 - ▶ Non-negative for any θ
 - ▶ Zero only if $q_{\theta}^{(0)}(z) = p(z | x)$

Variational Contrastive Divergence

- ▶ Key idea: The improved distribution $q_\theta(z)$ decreases the KL

$$\text{KL}(q_\theta(z) \parallel p(z \mid x)) \leq \text{KL}(q_\theta^{(0)}(z) \parallel p(z \mid x))$$

(equality only if $q_\theta^{(0)}(z) = p(z \mid x)$)

- ▶ A first objective:

$$\mathcal{L}(\theta) = \text{KL}(q_\theta^{(0)}(z) \parallel p(z \mid x)) - \text{KL}(q_\theta(z) \parallel p(z \mid x))$$

(it is a proper divergence)

Variational Contrastive Divergence

$$\mathcal{L}(\theta) = \text{KL}(q_\theta^{(0)}(z) \parallel p(z|x)) - \text{KL}(q_\theta(z) \parallel p(z|x))$$

- ▶ Still intractable: $\log q_\theta(z)$ in the second term
- ▶ Add regularizer,

$$\mathcal{L}_{\text{VCD}}(\theta) = \underbrace{\text{KL}(q_\theta^{(0)}(z) \parallel p(z|x)) - \text{KL}(q_\theta(z) \parallel p(z|x))}_{\geq 0} + \underbrace{\text{KL}(q_\theta(z) \parallel q_\theta^{(0)}(z))}_{\geq 0}$$

(still a proper divergence)

Variational Contrastive Divergence

$$\mathcal{L}_{\text{VCD}}(\theta) = \text{KL}(q_\theta^{(0)}(z) \parallel p(z|x)) - \text{KL}(q_\theta(z) \parallel p(z|x)) + \text{KL}(q_\theta(z) \parallel q_\theta^{(0)}(z))$$

- ▶ Addresses Challenge #1 (intractability):
 - ▶ The intractable term $\log q_\theta(z)$ cancels out
- ▶ Addresses Challenge #2 (weak dependence):
 - ▶ $\mathcal{L}_{\text{VCD}}(\theta) \xrightarrow{t \rightarrow \infty} \text{KL}(q_\theta^{(0)}(z) \parallel p(z|x)) + \text{KL}(p(z|x) \parallel q_\theta^{(0)}(z))$

Taking Gradients of the VCD

$$\mathcal{L}_{\text{VCD}}(\theta) = -\mathbb{E}_{q_\theta^{(0)}(z)} \left[\log p(x, z) - \log q_\theta^{(0)}(z) \right] + \mathbb{E}_{q_\theta(z)} \left[\log p(x, z) - \log q_\theta^{(0)}(z) \right]$$


- ▶ The first component is the (negative) standard ELBO
 - ▶ Use reparameterization or score-function gradients
- ▶ The second component is the new part,
$$\nabla_\theta \mathbb{E}_{q_\theta(z)} [g_\theta(z)] = -\mathbb{E}_{q_\theta(z)} \left[\nabla_\theta \log q_\theta^{(0)}(z) \right] + \mathbb{E}_{q_\theta^{(0)}(z_0)} \left[\mathbb{E}_{Q^{(t)}(z|z_0)} [g_\theta(z)] \nabla_\theta \log q_\theta^{(0)}(z_0) \right]$$
(can be approximated via Monte Carlo)

Algorithm to Optimize the VCD

$$\mathcal{L}_{\text{VCD}}(\theta) = -\mathbb{E}_{q_{\theta}^{(0)}(z)} \left[\log p(x, z) - \log q_{\theta}^{(0)}(z) \right] + \mathbb{E}_{q_{\theta}(z)} \left[\log p(x, z) - \log q_{\theta}^{(0)}(z) \right]$$

1. Sample $z_0 \sim q_{\theta}^{(0)}(z)$ (reparameterization)
2. Sample $z \sim Q^{(t)}(z | z_0)$ (run t MCMC steps)
3. Estimate the gradient $\nabla_{\theta} \mathcal{L}_{\text{VCD}}(\theta)$
4. Take gradient step w.r.t. θ

Toy Experiments



Optimizing the VCD leads to a distribution $q_{\theta}^{(0)}(z)$ with higher variance

$$\mathcal{L}_{\text{VCD}}(\theta) \xrightarrow{t \rightarrow \infty} \text{KL}_{\text{sym}}(q_{\theta}^{(0)}(z) \parallel p(z \mid x))$$

Experiments: Latent Variable Models

- ▶ Model is $p_\phi(x, z) = \prod_n p(z_n)p_\phi(x_n | z_n)$
- ▶ Amortized distribution $q_\theta(z_n | x_n) = \int Q^{(t)}(z_n | z_0) q_\theta^{(0)}(z_0 | x_n) dz_0$
- ▶ Goal: Find model parameters ϕ and variational parameters θ


method	average test log-likelihood	
	MNIST	Fashion-MNIST
Explicit + KL	-111.20	-127.43
Implicit + KL [Hoffman, 2017]	-103.61	-121.86
VCD	-101.26	-121.11

(a) Logistic matrix factorization

method	average test log-likelihood	
	MNIST	Fashion-MNIST
Explicit + KL	-98.46	-124.63
Implicit + KL [Hoffman, 2017]	-96.23	-117.74
VCD	-95.86	-117.65

(b) VAE

Summary



- ▶ Use *implicit distributions* to form expressive variational posteriors
 - Density ratio estimation
 - Semi-implicit distributions (SIVI, UIVI)
 - Refine the variational distribution with MCMC (VCD)
- ▶ Stable training
- ▶ Good empirical results on (deep) probabilistic models

Proof of the Key Equation in UIVI

- Goal: Prove that

$$\nabla_z \log q_\theta(z) = \mathbb{E}_{q_\theta(\varepsilon | z)} [\nabla_z \log q_\theta(z | \varepsilon)]$$

- Start with log-derivative identity,

$$\nabla_z \log q_\theta(z) = \frac{1}{q_\theta(z)} \nabla_z q_\theta(z)$$

- Apply the definition of $q_\theta(z)$ through a mixture,

$$\nabla_z \log q_\theta(z) = \frac{1}{q_\theta(z)} \int \nabla_z q_\theta(z | \varepsilon) q(\varepsilon) d\varepsilon$$


- Apply the log-derivative identity on $q_\theta(z | \varepsilon)$,

$$\nabla_z \log q_\theta(z) = \frac{1}{q_\theta(z)} \int q_\theta(z | \varepsilon) q(\varepsilon) \nabla_z \log q_\theta(z | \varepsilon) d\varepsilon.$$

- Apply Bayes' theorem

UIVI Experiments: Multinomial Logistic Regression

$$p(x, z) = p(z) \prod_{n=1}^N \frac{\exp\{x_n^\top z_{y_n} + z_{0y_n}\}}{\sum_k \exp\{x_n^\top z_k + z_{0k}\}}$$



Number of HMC iterations does not significantly impact results

Generalized VCD

- ▶ VCD

$$\mathcal{L}_{\text{VCD}}(\theta) = \text{KL}(q_{\theta}^{(0)}(z) \parallel p(z \mid x)) + \text{KL}(q_{\theta}(z) \parallel q_{\theta}^{(0)}(z)) - \text{KL}(q_{\theta}(z) \parallel p(z \mid x))$$

- ▶ α -generalized VCD ($0 < \alpha \leq 1$)

$$\mathcal{L}_{\text{VCD}}^{(\alpha)}(\theta) = \text{KL}(q_{\theta}^{(0)}(z) \parallel p(z \mid x)) + \alpha \left[\text{KL}(q_{\theta}(z) \parallel q_{\theta}^{(0)}(z)) - \text{KL}(q_{\theta}(z) \parallel p(z \mid x)) \right]$$

VCD Experiments: Impact of Number of MCMC Steps

